

Supplementary Table 1. Patient and tumor factors predicting local tumor control

Factors	Univariable		Multivariable	
	<i>p</i> -value	OR (95% CI)	<i>p</i> -value	OR (95% CI)
Age	0.23	0.97 (0.92-1.01)	0.17	0.94 (0.87-1.02)
WHO performance	0.88	1.08 (0.38-3.15)	0.13	5.60 (0.61-59.30)
Fitness for surgery	0.54	0.71 (0.25-2.08)	0.32	0.36 (0.05-2.76)
Initial T stage	0.48	0.39 (0.03-5.21)	0.69	0.51 (0.02-14.10)
Initial N stage	0.67	1.38 (0.31-6.14)	0.69	0.64 (0.07-5.76)
Distance from anal verge	0.94	1.05 (0.34-3.21)	0.55	0.63 (0.14-2.86)
Tumor size	0.47	0.50 (0.78-3.22)	1.00	0.00 (0.00-0.00)
EBRT regimen	0.94	0.95 (0.25-3.60)	0.63	0.56 (0.06-5.70)
Clinical response after EBRT	0.81	4.30 (0.84-22.20)	0.28	2.93 (0.41-20.85)
yT-stage at the time of CXB	0.12	3.67 (0.70-19.17)	0.57	1.95 (0.20-19.57)
CXB total dose	0.73	0.69 (0.08-5.64)	0.97	1.05 (0.06-18.01)